

Passengers' destinations from China: Low risk of Novel Coronavirus (2019-nCoV) transmission into Africa and South America

NOTE: The following is an accepted manuscript for *Epidemiology and Infection*. Once published the paper will be available at:
<https://doi.org/10.1017/S0950268820000424>

Najmul Haider¹, Alexei Yavlinsky², David Simons¹, Abdinasir Yusuf Osman¹, Francine Ntoumi^{3,4}, Alimuddin Zumla⁵, Richard Kock¹

¹ The Royal Veterinary College, University of London, Hawkshead Lane, North Mymms, Hatfield, Hertfordshire

²Institute of Health Informatics, University College London, United Kingdom

³Fondation Congolaise pour la Recherche Médicale (FCRM), Brazzaville, Republic of Congo.

⁴Institute for Tropical Medicine, University of Tübingen, Germany.

⁵UCL Centre for Clinical Microbiology, Department of Infection, Division of Infection and Immunity, Royal Free campus, and NIHR Biomedical Research Centre, UCL Hospitals NHS Foundation Trust, London, UK, London, United Kingdom

Abstract:

Novel Coronavirus (2019-nCoV) was detected in humans during last week of December 2019 at Wuhan city in China, and caused 24,554 cases in 27 countries and territories as of February 5th 2020. The objective of this study was to estimate the risk of transmission of 2019-nCoV through human passenger air flight from four major cities of China (Wuhan, Beijing, Shanghai, and Guangzhou) to the passengers' destination countries.

We extracted the weekly simulated passengers' end destination data for the period of 1st to 31st January 2020 from FLIRT, an online air travel dataset that uses

information from 800 airlines to show the direct flight and passengers' end destination. We estimated a risk index of 2019-nCoV transmission based on the number of travellers to destination countries, weighted by number of confirmed cases of the departed city reported by WHO. We ranked each country based on risk index in four quantiles (4th quantile being the highest risk and 1st quantile being the lowest risk).

During the period 388,287 passengers were destined for 168 countries or territories across the world. The risk index of 2019-nCoV among the countries had very high correlation with WHO reported confirmed cases (0.97). According to our risk score classification, of the countries that reported at least one 2019-nCoV case as of February 5th, 2020, 24 countries were in the fourth quantile of the risk index, two in the third quantile, one in the second quantile and none in the first quantile. Outside China, countries with higher risk of 2019-nCoV transmission are Thailand, Cambodia, Malaysia, Canada and United States of America (USA), all of which reported at least one case. In pan-Europe, United Kingdom, France, Russia, Germany and Italy, in North America, USA and Canada, in Oceania, Australia had high risk, all of them reported at least one case. In Africa and South America, the risk of transmission is very low with Ethiopia, South Africa, Egypt, Mauritius and Brazil showing similar risk of transmission compared to the risk of any of the countries where at least one case is detected.

The risk of transmission at 31st January 2020 was very high in neighbouring Asian countries, followed by Europe (UK, France, Russia, and Germany), Oceania (Australia, New Zealand and Belize) and North America (USA and Canada).

Increased public health response including early case recognition, isolation of identified case, contact tracing and targeted airport screening, public awareness and vigilance of health workers will help mitigate the force of further spread to naïve countries.

Key words: 2019-nCoV, China, Wuhan, Risk map, Africa, transmission

Introduction:

On the 31st of December 2019 local hospitals in Wuhan, China reported that they had detected a series of cases of novel coronavirus-infected pneumonia to World Health Organization (WHO) [1]. On January 7th the causative agent was identified by the Chinese Centre for Disease Control and Prevention as a novel coronavirus and designated “2019-nCoV”. Epidemiological investigations identified the local Huanan seafood wet market as the location of an initial exposure event [2]. The market was closed on the 31st of December 2019 [2,3] and wildlife market activity banned countrywide. Despite travel restrictions to and from the city imposed by Chinese authorities to limit the potential dispersion of the virus beyond the region, [4,5] international cases continue to be reported.

As of February 5th 2020 there were 24,554 confirmed cases in 27 countries or territories, of which 24363 (99.2%) were within mainland China[6]. The locations of internationally imported cases are consistent with risk models generated from flight data out of Wuhan city. Transmission from mildly symptomatic (i.e. cough, lethargy, myalgia) infected individuals was identified early in the course of this outbreak, with human-to-human transmission detected in international case series [7].

The timing of this outbreak around the lunar new year widely celebrated in China coincides with an annual period of high annual human movement patterns in the region and between China and the rest of the world [8], increasing the potential for rapid geographic dispersal of the infection. Further, recent investment in the African continent by the Chinese state and private investors has led to an increasing Chinese diaspora [9] and a greater number of direct and indirect flight connections to the African continent from China [10].

There are few studies available on global risk of 2019-nCoV spread [4,5,11]. Bogoch et al.[5] and Chinazzi et al.[11] estimated the risk of importation of 2019-nCoV from major Chinese cities to the most frequent international destinations. Wu et al. estimated the risk of international spread compared to domestic outbound flights [4]. These articles do not model the cumulative risk of importation of 2019-nCoV in a country and instead focus on specific points of entry. Here, we considered all the end destinations of flights from four important cities of China involving 168 countries/territories around the world and calculated the total risk for transmission into a country by aggregating risk associated with all the entry airports of the country. We further looked in more detail at the risk to Africa where the health infrastructure would be challenged tracking a new epidemic across its 54 countries.

The aim of the current study was to explore the effect of sustained transmission from the four Chinese cities of Wuhan, Beijing, Shanghai and Guangzhou on international disease importation risk to 168 countries and territories, with a specific focus on Africa where current levels of healthcare infrastructure could provide a significant challenge for managing this novel epidemic.

Methods:**Data:**

We extracted modelled flight data for the final destination of passengers travelling between from four Chinese cities (including domestic and international destinations) from the FLIRT database [12,13] . FLIRT was designed to predict the flow and likely destination of infected travellers through the air travel network. It uses a database of flight schedules from over 800 airlines it displays direct flight connections in addition to a modelled end destination (3 letter IATA code for airports). Flight connection data and passenger numbers are based on data collected since October, 2014[12]. We extracted the simulated passenger's data for each week for the period of 1st January to 31st January 2020 for four major Chinese cities: Wuhan, Beijing, Shanghai, and Guangzhou. The simulation can process up-to 20,000 passengers' information for a particular time frame from any city (including surrounding airports). We collected reported 2019-nCoV case data from WHO's daily situation update website [14].

Estimation of risk of transmission:

To estimate the relative risk of 2019-nCoV transmission we considered all infected passengers who travelled between 1st January and 31st January to possess a maximum risk of transmission 1 (and no infected passengers means no risk) and estimated the relative risk of each country based on the number of passengers which travelled from each of the four cities. Thus countries with a higher number of passengers travelling from any of these cities had higher risk of transmission. We then weighted the risk estimated for each city with the number of reported infected

people in each city by 31st January 2020 [14] and estimated mean average risk of transmission termed as “Risk index” which follows the equation below.

$$Risk\ index(x) = \sum_{n=1}^N \frac{P(x)_n}{P_n} \times \frac{I_n}{C_n}$$

Where x is the destination country, $Risk\ index(x)$ is the risk of infection importation in country x , $P(x)_n$ is the number of passengers to country x from city n , P_n is the total number of passengers who left city n , I_n is the number of infected people in city n and C_n is the population size of city n . The risk index denotes the risk of at least one case being imported into a country or territory where 1 means an absolute certainty and 0 means no risk at all. Our model assumed that there is no case outside China and thus ignored if any country already had imported case(s). In countries, where 2019-nCoV is already detected, the risk index would be explain the risk of importing additional infected individuals from China.

We performed a Pearson correlation coefficient test between the risk index of the country and the WHO’s reported case number from the country. We grouped the countries in four quantiles based on the risk index where high-risk countries were grouped as the 4th (>75th percentiles) and the 3rd (>50th percentiles ≤75th percentiles) quantiles and low-risk countries were grouped as the 2nd (>25th percentiles ≤50th percentiles) and the 1st (≤25th percentile) quantiles (Table 1).

Results:

We modelled 388,287 passengers travelling to 168 countries or territories. The risk index of 2019-nCoV for these countries are presented in figure 1. A regularly updated risk map is hosted on PANDORA's website (<https://ncovdata.io/import/>).

Outside China the countries with the highest risk of 2019-nCoV transmission from our model were Thailand, Cambodia, Malaysia, Canada and the United States of America (USA), all of which have reported at least one case. Among the top 25 countries identified with the highest risk of 2019-nCoV transmission (Fig 2), all countries except four (Indonesia, Turkey, Pakistan and Qatar) have detected at least one case as of February 5th 2020 (Table 1).

According to our risk score classification, of the countries that reported at least one 2019-nCoV case as of February 5th 2020, 24 countries were in the fourth quantile of the risk index, two (Sweden and Belgium) in the third quantile, one (Nepal) in the second quantile and none in the first quantile [14]. Asian and European countries are dominant in the 3rd and 4th quantile (high risk index) while African and South American Countries are the majority in the 1st and 2nd quantiles (low risk index) (Table 1). Out of 43 countries in the 4th quantile, 22 were from Asia and 13 from Pan-Europe, whereas in the 1st quantile 19 out of 42 countries were from Africa. (Table 1)

The overall risk of transmission of the virus into Africa is low. However, Ethiopia, South Africa, Egypt and Mauritius have a similar risk score as countries where at least one case was detected. In South America, only Brazil has similar or greater risk than countries currently reporting cases. In North America, both USA and Canada have high risk and had imported cases reported early in the outbreak. Australia and New Zealand have risk similar to the countries where at least one case is detected.

Although there are few direct flights from China to African destinations, a large number of indirect flights operate via Dubai, an international airport hub in the United Arab Emirates.

The correlation coefficient between estimated risk index of the countries and WHO reported confirmed cases was 0.97.

Discussion:

Our analysis showed a high risk of transmission of 2019-nCoV through air flights from four Chinese cities to neighbouring Asian countries. The risk of 2019-nCoV transmission was relatively low in Africa and South America. Several countries in both North America and Oceania showed high risk with these countries reporting at least one case of 2019-nCoV. Our risk index showed a very high correlation with WHO's reported 2019-nCoV cases.

China has four times as many air passengers now than it had during SARS outbreaks in 2003. A large number of workers now travel internationally where China is heavily investing in infrastructure development in Africa, parts of Asia and Latin

America. A significant and mobile Chinese population live in Europe and North America alongside an increasing amount of Chinese tourism. This travel poses a high risk of 2019-nCoV travelling across international borders. Although the acquiring a case is low for these countries, the consequences are likely to be higher because of the country's capacity to control such situations[15].

Based on our model the countries with the highest risk index but have not reported any case of 2019-nCoV as yet are Indonesia, Pakistan, Turkey, Qatar and Ethiopia. These countries are at risk and they should be priorities for investment in case detection and airport screening. Compared to the SARS outbreak of 2004, the situation in 2020 differs due to the increased frequency and volume of international air travel.

During these early stages of the epidemic case numbers have doubled on average every 7.4 days with an estimated basic reproduction number (R_0) of 2.2 (1.4-3.9)[1]. Although the data so far suggests that the disease is mild in most cases and that the case fatality rate is currently reported to be lower than SARS or MERS, the situation is likely to go on for months and could cause severe disruption in countries that are not well prepared. Hence countries ranked as high risk in our model (4th and 3rd quantiles) should take all steps necessary to ensure prompt detection of cases and capacity to manage these cases to prevent ongoing spread. International investment needs to be directed especially to countries with limited healthcare and public health surveillance capacity to enable detection of cases and disease control [16,17]

Our estimation showed a lower risk of transmission in Africa and South America. Nevertheless, low and middle countries on these continents are more likely to see ongoing spread and major disruption from the introduction of a single case, even if the risk of importation is lower. Direct flights between Chinese cities and African countries are few which has contributed to a lower estimated risk of 2019-nCoV transmission. As of 5th February 2020, five cases have been reported from the United Arab Emirates (UAE) which acts as an important travel hub for onward journeys to the African continent. Implementation of mildly symptomatic passenger screening in the UAE may reduce the potential for 2019-nCoV to enter Africa. Screening and diagnostic capacity in Africa has been supported by a rapid grant from the Bill and Melinda Gates Foundation to the African CDC with the aim of mitigating the consequences of an importation of 2019-nCoV case.

The current situation is an extremely dynamic and since then some countries have instigated flight restrictions and closed borders (e.g. Russia). These decisions were relevant for these locations but not based on probabilities. WHO has not recommended a cessation of transportation to free countries but suggested preventive measures. This would seem appropriate for Africa and South America with the caveat that only one case is needed to initiate a local epidemic without proper biosecurity and quarantine measures, whilst other regions will need to decide on a case-by-case basis through appropriate risk assessment.

Our study has several limitations. We considered flights from four cities of China, three of which (Beijing, Shanghai, and Guangzhou) are ranked among top five busiest airports (based on number of flights) in China and Wuhan as the site of origin

of outbreaks. While including further cities in our analysis would have added further information, Beijing and Shanghai cover most of the international destinations to which other airports are connected. Further, we have adjusted for the number of reported cases in each departure. When developing the model we initially explored using only Wuhan as the departure airport, the rank of top 10 at risk countries remained the same. Thus our findings is still representative of total risk posed by other airports or cities. We did not consider the risk associated with the travel by water and land which might have an impact in spread of 2019-nCoV. Another limitation is that the model does not account for travel patterns in other affected countries. For example some cases have started acquiring the disease outside of China: the third case notified in the UK acquired the disease in Singapore. However overall the risk of this compared to the risk of acquisition in China is very low and would therefore have minimal impact on the the order of risk of the countries presented here.

Conclusion:

The risk of transmission from China at January 31st 2020 was highest to neighbouring countries in Asia (Thailand, Cambodia, Malaysia), followed by Europe (UK, France, Russia, and Germany), Oceania (Australia, and New Zealand) and North America (USA and Canada). The situation is dynamic and may have changed with the closure of flights and borders since this analysis was done. The higher correlation coefficient with travellers and case detection data indicate that 2019-nCoV will remain a significant threat from air -borne movement of people. The authors suggest an ongoing risk based approach to prioritisation of and investment by international and national agencies and authorities, in emergency interventions for the prevention of movement of the 2019_nCoV through human travel. This could be

achieved by appropriate actions at high-risk points of departure and at highly used ports of entry from these infected zones. Closure of certain routes, targeted airport screening, risk communication, public awareness and targeted training and vigilance of health workers associated with portals of entry of visitors to their countries will help mitigate the force of further spread of 2019_nCoV.

Acknowledgements: All authors are part of PANDORA-ID-NET Consortium (EDCTP Reg/Grant RIA2016E-1609) funded by the European and Developing Countries Clinical Trials Partnership (EDCTP2) programme which is supported under Horizon 2020, the European Union's Framework Programme for Research and Innovation. AZ is in receipt of an National Institutes of Health Research (NIHR) senior investigator award and the PANDORA-ID-NET. FN and AZ acknowledge support from EDCTP (CANTAM2). We acknowledge Eco Health Alliance for making FLIRT data publicly available.

Conflict of interest: The authors declare that they have no conflict of interest

References:

1. **Li Q, et al.** Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia. *The New England journal of medicine* 2020; : 1–9.
2. **Chen N, et al.** Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet (London, England)* Elsevier Ltd, 2020; **6736**: 1–7.
3. **ProMED.** *Undiagnosed pneumonia - China (HU) (01): wildlife sales, market closed, RFI.* 2020.
4. **Wu JT, Leung K, Leung GM.** Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *The Lancet* Elsevier Ltd, 2020; **6736**Published online: January

- 2020.doi:10.1016/S0140-6736(20)30260-9.
5. **Bogoch II, et al.** Potential for global spread of a novel coronavirus from China. *Journal of Travel Medicine* 2020; Published online: 2020.
 6. **Johns Hopkins University.** *Coronavirus 2019-nCoV.*
 7. **Rothe C, et al.** Transmission of 2019-nCoV Infection from an Asymptomatic Contact in Germany. *The New England journal of medicine* 2020; : 2019–2020.
 8. **Wang X, et al.** Tracing The Largest Seasonal Migration on Earth. *arXiv* 2014; Published online: November 2014.
 9. **Brautigam D, Xiaoyang T, Xia Y.** What kinds of Chinese ‘geese’ are flying to Africa? Evidence from Chinese manufacturing firms. *Journal of African Economies* 2018; Published online: 2018.doi:10.1093/jae/ejy013.
 10. **Wang J, Yang H, Wang H.** The evolution of China’s international aviation markets from a policy perspective on air passenger flows. *Sustainability (Switzerland)* 2019; **11**: 3566.
 11. **Chinazzi M, et al.** Preliminary assessment of the International Spreading Risk Associated with the 2019 novel Coronavirus (2019-nCoV) outbreak in Wuhan City. 2020; : 1–7.
 12. **Huff A, et al.** FLIRT-ing with Zika: A Web Application to Predict the Movement of Infected Travelers Validated Against the Current Zika Virus Epidemic. *PLoS Currents* 2016.
 13. **Alliance EH.** *FLIRT: A Product of EcoHealth Alliance.* Available at: <https://flirt.eha.io/>. 2016.
 14. **WHO.** *Novel Coronavirus(2019-nCoV) Situation Report - 1-14.* 2020.
 15. **Johns Hopkins Center for Health Security.** *Global Health Security Index.* 2020.
 16. **Hui DS, et al.** *The continuing 2019-nCoV epidemic threat of novel coronaviruses to global health — The latest 2019 novel coronavirus outbreak in Wuhan, China.* *International Journal of Infectious Diseases.* 2020.
 17. **Mccloskey B, Heymann DL.** SARS to novel coronavirus – old lessons and new lessons. *Epidemiology and Infection* 2020; **148**: 3–6.

Figures and Tables:

Figure legends:

Fig 1: The map with risk index of countries with 2019-nCoV infection (0-1). The darker colour indicate higher risk and light blue colour indicate absence of data. In general, China and neighbouring countries has higher risk of transmission of 2019-nCoV infection. Africa and South America generally have a low risk of transmission. Ethiopia, South Africa, Egypt, Mauritius and Brazil have similar risk of transmission to countries where at least one case has been detected. For example, the risk index of 0.1 for Thailand indicates that based on travel patterns observed during January 1st -31st 2020 from four major cities of China, Thailand has 10% risk of importing a 2019-nCoV infected person from China.

Fig 2: Chart showing relative risk of countries outside China being exposed to coronavirus (2019-nCoV) transmission. The second Y axis indicates number of confirmed 2019-nCoV cases reported by WHO as of 5th February 2020. Twenty one of the top 25 at-risk countries (except Indonesia, Turkey, Pakistan, and Qatar) reported at least one 2019-nCoV case by 5th February 2020.

Fig 1:




Fig 2:




Table 1: The list of Countries based with their risk index in different quantiles for 2019-nCoV transmission

SI/Rank	4 th Quantile risk index (highest risk)		3 rd Quantile risk index		2 nd Quantile risk index		1 st Quantile risk index (lowest risk)	
	Country	Risk index	Country	Risk index	Country	Risk index	Country	Risk index
1	China	0.609603126	Sweden	0.000320248	Kenya	3.53E-05	Jamaica	3.84E-06
2	Thailand	0.099432816	Laos	0.000296262	Peru	3.18E-05	Serbia	3.22E-06
3	Cambodia	0.05294058	Brazil	0.00027186	Algeria	3.11E-05	Togo	2.73E-06
4	Malaysia	0.041899039	Denmark	0.000254904	French Polynesia	3.07E-05	Uganda	2.71E-06
5	Canada	0.02730388	Oman	0.000248884	Iceland	3.00E-05	Tonga	2.69E-06
6	United States	0.021169936	Israel	0.000221865	Samoa	2.74E-05	The Bahamas	2.39E-06
7	Japan	0.01479856	Ukraine	0.000209515	Tanzania	2.73E-05	Cote d'Ivoire	2.02E-06
8	India	0.010256629	Poland	0.0002	Palau	2.63E-05	Suriname	2.01E-06
9	United Kingdom	0.008786839	Brunei	0.000181028	Djibouti	2.45E-05	Vanuatu	1.99E-06
10	South Korea	0.008072566	Czech Republic	0.000179806	Belarus	2.40E-05	Albania	1.90E-06
11	Vietnam	0.007928803	Northern Mariana Islands	0.000177972	Bosnia and Herzegovina	2.40E-05	Malta	1.73E-06
12	Singapore	0.007784474	Belgium	0.000175566	Cook Islands	2.26E-05	Guinea	1.60E-06

13	Hong Kong	0.007636714	Maldives	0.000172453	Colombia	1.94E-05	Namibia	1.55E-06
14	Indonesia	0.007197131	Norway	0.000171441	Papua New Guinea	1.88E-05	Democratic Republic of Congo	1.41E-06
15	United Arab Emirates	0.007089607	Kuwait	0.000166929	Nigeria	1.85E-05	Rwanda	1.22E-06
16	France	0.006814962	Egypt	0.000139373	Jordan	1.83E-05	Honduras	1.02E-06
17	Turkey	0.00568808	Iran	0.000137181	Cuba	1.79E-05	Gabon	8.98E-07
18	Australia	0.005671745	Mongolia	0.000132319	Argentina	1.70E-05	Republic of Congo	7.22E-07
19	Russia	0.005592859	Chile	0.000129975	Tunisia	1.65E-05	Bermuda	5.99E-07
20	Pakistan	0.004811284	North Korea	0.000128562	Ghana	1.61E-05	Antigua and Barbuda	3.52E-07
21	Qatar	0.004113225	Mauritius	0.000126679	Armenia	1.53E-05	Barbados	3.52E-07
22	Macau	0.003373727	Portugal	0.000113251	Dominican Republic	1.46E-05	Cape Verde	3.52E-07
23	Germany	0.003176858	Uzbekistan	9.67E-05	New Caledonia	1.34E-05	Guyana	3.52E-07
24	Italy	0.002753413	Hungary	8.97E-05	Cyprus	1.29E-05	Madagascar	2.99E-07
25	Philippines	0.002740638	Azerbaijan	8.64E-05	Bhutan	1.25E-05	Grenada	2.47E-07
26	Taiwan	0.002590034	Croatia	8.57E-05	Nepal	1.25E-05	Bolivia	1.76E-07
27	Belize	0.002009996	Tajikistan	8.50E-05	Slovenia	1.24E-05	Burkina Faso	1.76E-07
28	Ethiopia	0.001469205	Bahrain	6.31E-05	Moldova	1.22E-05	Cameroon	1.76E-07
29	Finland	0.001307074	Fiji	6.29E-05	Kosovo	1.21E-05	Chad	1.76E-07
30	Sri Lanka	0.001179859	Kyrgyzstan	6.22E-05	Zambia	1.20E-05	Mozambique	1.76E-07
31	Netherlands	0.000980799	Afghanistan	6.16E-05	El Salvador	1.05E-05	Paraguay	1.76E-07
32	New Zealand	0.000971254	Panama	5.91E-05	Romania	7.56E-06	Solomon Islands	1.76E-07

33	Greece	0.000958209	Morocco	5.60E-05	Guatemala	6.92E-06	Syria	1.76E-07
34	Bangladesh	0.000831196	Iraq	5.54E-05	Angola	6.77E-06	Uruguay	1.76E-07
35	Myanmar	0.000803755	Bulgaria	5.07E-05	Costa Rica	6.41E-06	Venezuela	1.76E-07
36	Saudi Arabia	0.000750981	Lithuania	4.99E-05	Trinidad and Tobago	6.13E-06	Zimbabwe	1.76E-07
37	Spain	0.000564876	Seychelles	4.82E-05	Sudan	5.66E-06	Libya	1.23E-07
38	Switzerland	0.00056232	Lebanon	4.47E-05	Ecuador	5.56E-06	Macedonia	1.23E-07
39	Austria	0.000498664	Georgia	4.37E-05	Puerto Rico	4.68E-06	Saint Lucia	1.23E-07
40	South Africa	0.000468876	Latvia	4.06E-05	Turkmenistan	4.42E-06	Sierra Leone	1.23E-07
41	Kazakhstan	0.000443175	Luxembourg	3.80E-05	Mauritania	4.09E-06	Somalia	1.23E-07
42	Ireland	0.000371199	Estonia	3.69E-05			Timor-Leste	1.23E-07
43	Mexico	0.000322198						

44

Number of countries/
territories: 168

Africa : 2
Asian : 22
Pan-Europe :13
North America: 4
Oceania: 2
South America: 0
Total: 43

Africa : 4
Asian : 16
Pan-Europe : 17
North America: 0
Oceania: 2
South America: 3
Total: 42

Africa : 11
Asian : 4
Pan-Europe : 9
North America: 3
Oceania: 6
South America: 8
Total: 41

Africa : 19
Asian : 2
Pan-Europe : 4
North America: 9
Oceania: 3
South America: 7
Total: 42